



Pseudovirus Ebola Zaire (luciferase)

Lot #PsV-EBZLU-0100



Quality control report

1. Summary

The lot number #PsV-EBZLU-0100 is a lentivirus-based pseudovirus pseudotyped with the glycoprotein of the Ebola Zaire variant. This quality control report demonstrates that the lot #PsV-EBZLU-0100 is efficient for cell transduction and can be effectively neutralized by a standard neutralizing antibody.

2. Transduction efficiency assay

Target cells	HEK293 cells (TIM-1 ⁺)
Volume of pseudovirus	0, 1, 5, 10, 20, 30 or 40 μ L/well
Detection signal	Luminescence (firefly luciferase)
Detection method	Microplate reader




Figure 1: Transduction efficiency curve.

A volume range of pseudovirus was mixed in a final volume of 50 μ L of culture complete medium, in a 96-well plate. Then, an additional 50 μ L containing 20 000 HEK293 cells (TIM-1⁺) was seeded in each well. Luc expression analysis was performed 48 hours post-infection by a luminescence microplate reader.

Conclusion	The Luc Ebola Zaire pseudovirus (#PsV-EBZLU-0100) can transduce the target cells.
-------------------	---

3. Neutralization assay

Target cells	HEK293 cells (TIM-1 ⁺)
Volume of pseudovirus	1 μ L/well
Neutralizing antibody (Nabs)	Anti-Ebola surface glycoprotein [KZ52], AB00690-10.0
Detection signal	Luminescence (firefly luciferase)
Detection method	Microplate reader




Figure 2: Neutralization curve.

A monoclonal neutralizing antibody ([AB00690-10.0](#)) at 1 μ g/ μ L was serially diluted in a final volume of 50 μ L of complete medium and incubated for 1 hour at 37 °C, with 1 μ L of pseudovirus, in a 96-well plate. Then, an additional 50 μ L containing 20 000 HEK293 cells (TIM-1⁺) was seeded in each well and incubated for 48 hours. Finally, an additional 100 μ L of [Bright-Glo™ Luciferase](#) buffer was added in each well and incubated for 2 minutes. Data in relative unit luminescence (RLU) were obtained from the analysis of 150 μ L of the cell lysate with a microplate reader. Raw data were analyzed using a log(inhibitor) vs normalized-response (variable slope) non-linear regression model in Prism v10 (GraphPad). Percentages of neutralization were normalized considering only cells into wells as 100% neutralization and cells transduced by pseudoviruses without any NABs as 0% neutralization. Data are representative of duplicates.

Conclusion

The Luc Ebola Zaire pseudovirus #PsV-EBZLU-0100 can be efficiently neutralized by neutralizing antibodies.

3. Additional information

Caution

We recommend determining the optimal pseudovirus volume to use according to your specific experimental conditions.

Pseudovirus

3rd generation, replication incompetent

Pseudotyping

Ebola virus glycoprotein from the 2014 outbreak in west Africa (GENBANK : [KP096420.1](#))

Glycosylation origin

Human

Reporter protein

Firefly luciferase

Storage

- 80 °C, avoid freeze/thaw cycles

For more information

mathias.mangion@ivanobioscience.com

Message object should contain : "#PsV-EBZLU-0100"